



STEVEN C. KRONLAGE, MD
Hand & Upper Extremity Surgery
Florida Bone & Joint Specialists



POST OP INSTRUCTIONS – DEQUERVAIN’S RELEASE/GANGLION CYST

- **Keep hand and arm elevated most of the time.**
- The numbness from surgery usually lasts about 6 hours.
- You will have a coban dressing and a removable splint. The splint can be readjusted and even removed but should be on when you are doing even light activities, going out, and sleeping.
- On post op day _____ it is OK to remove the bandages entirely. You can wash with soapy water, but do not soak or immerse in water (no dishes, hot tubs, etc).
- I want you to move your fingers freely and actively (full range of motion), just don’t lift anything more than a pound or two.
- If you had surgery on your dominant hand, writing will be difficult. Don’t push it.
- The splint is a reminder to not lift anything. Wear it when you are out and about and to sleep.
- Do not use over the counter salves, Neosporin, Bacitracin or other ointments on the wound. Hands and fingers sweat and the wound will get macerated and open up.
- Band aids are OK, but make sure to let the incision “breathe” as it will heal much faster with air.
- Call us to be evaluated for any redness, increased heat or anything that looks like an infection.

850.916.8480

1040 Gulf Breeze Parkway, Suite 209; Gulf Breeze, FL 32561
Phone: 850.916.8480 Fax: 850.916.8499 Web: www.stevenkronlage.com